

Electron energy spectrum in 2D quantum dot

ŚLAWOMIR WOLSKI¹, VITALII DUGAEV¹, JÓZEF BARNAŚ², EVGENY SHERMAN^{3,4},
AND MICHAŁ INGLÓT¹

¹*Department of Physics and Medical Engineering, Rzeszów University of Technology,
Poland*

wolski@prz.edu.pl

²*Faculty of Physics, Adam Mickiewicz University, ul. Umultowska 85, 61-614
Poznań, Poland*

³*Department of Physical Chemistry, The University of the Basque Country
UPV/EHU, Bilbao, Spain*

⁴*IKERBASQUE Basque Foundation for Science, Bilbao, Spain*

ABSTRACT

Semiconductor quantum dots introduced into the photoelectric material are currently a very important and effective way to increase the efficiency of photoelectric devices and photovoltaic cells. This is related to the appearance of additional energy levels in the system, and it expands the range of absorption frequencies toward the infrared.

This work contains the theory and numerical calculation of energy levels in quantum dots in perovskite solids. The model of the structure includes a semiconductor quantum dot shaped in nanodisk which is surrounded by another semiconductor. The calculations of the energy levels as functions of the dot radius and various types of semiconductors materials are performed.

This work is supported by the National Center of Research and Development in Poland, project POIR.01.01.01-00-598/15.